

0760-0305P

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

10/089695 NEW

**INTERNATIONAL APPLICATION NO.**

**INTERNATIONAL FILING DATE**

PCT/JP00/05214

August 3, 2000

**PRIORITY DATE CLAIMED**

**TITLE OF INVENTION**

# METHOD FOR PROMOTING EFFICIENCY OF GENE TRANSFER INTO PLANT CELLS

**APPLICANT(S) FOR DO/EO/US**

HIEI, Yukoh; KASAOKA, Keisuke; ISHIDA, Yuji

**Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:**

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau. WO 02/12521
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).  
☒ a. ☒ is transmitted herewith.  
b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4)
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

**Items 11. to 20. below concern document(s) or information included:**

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98, Form PTO-1449(s), and International Search Report (PCT/ISA/210) with 0 cited document(s).
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
14. ☐ A SECOND or SUBSEQUENT preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information:
- ✓ 1.) PCT/IB/308
- ✓ 2.) Four (4) sheets of Formal Drawings

JC13 Rec'd PCT/PTO 03 APR 2002

U.S. APPLICATION NO (if known, see 37 CFR 1.5) <b>NEW 10/089695</b>		INTERNATIONAL APPLICATION NO PCT/JP00/05214		ATTORNEY'S DOCKET NUMBER 0760-0305P	
--	--	--	--	--	--

<p>21. <input checked="" type="checkbox"/> The following fees are submitted:</p> <p><b>BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5):</b>          Neither international preliminary examination fee (37 CFR 1.482)          nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO          and International Search Report not prepared by the EPO or JPO. .... <b>\$1,040.00</b></p> <p>International preliminary examination fee (37 CFR 1.482) not paid to          USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$890.00</b></p> <p>International preliminary examination fee (37 CFR 1.482) not paid to USPTO          but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. .... <b>\$740.00</b></p> <p>International preliminary examination fee (37 CFR 1.482) paid to USPTO          but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$710.00</b></p> <p>International preliminary examination fee (37 CFR 1.482) paid to USPTO          and all claims satisfied provisions of PCT Article 33(1)-(4) ..... <b>\$100.00</b></p> <p><b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b></p> <p>Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input checked="" type="checkbox"/> 20 <input type="checkbox"/> 30          months from the earliest claimed priority date (37 CFR 1.492(e)).</p> <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="width:20%;">CLAIMS</th> <th style="width:20%;">NUMBER FILED</th> <th style="width:20%;">NUMBER EXTRA</th> <th style="width:20%;">RATE</th> <th style="width:20%;"></th> <th style="width:20%;"></th> </tr> <tr> <td>Total Claims</td> <td>48 - 20 =</td> <td>28</td> <td>X \$18.00</td> <td>\$</td> <td>504.00</td> </tr> <tr> <td>Independent Claims</td> <td>1 - 3 =</td> <td>0</td> <td>X \$84.00</td> <td>\$</td> <td>0</td> </tr> <tr> <td colspan="3">MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes</td> <td>+ \$280.00</td> <td>\$</td> <td>280.00</td> </tr> <tr> <td colspan="4" style="text-align: right;"><b>TOTAL OF ABOVE CALCULATIONS =</b></td> <td>\$</td> <td>1804.00</td> </tr> </table> <p><input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are          reduced by 1/2.</p> <p style="text-align: right;"><b>SUBTOTAL =</b> \$ <b>1804.00</b></p> <p>Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30          months from the earliest claimed priority date (37 CFR 1.492(f)).</p> <p style="text-align: right;"><b>TOTAL NATIONAL FEE =</b> \$ <b>1804.00</b></p> <p>Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be          accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property +</p> <p style="text-align: right;"><b>TOTAL FEES ENCLOSED =</b> \$ <b>1804.00</b></p> <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:60%;"></td> <td style="width:20%; text-align: right;">Amount to be:</td> <td style="width:20%;"></td> </tr> <tr> <td></td> <td style="text-align: right;">refunded</td> <td>\$</td> </tr> <tr> <td></td> <td style="text-align: right;">charged</td> <td>\$</td> </tr> </table>	CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE			Total Claims	48 - 20 =	28	X \$18.00	\$	504.00	Independent Claims	1 - 3 =	0	X \$84.00	\$	0	MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes			+ \$280.00	\$	280.00	<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	1804.00		Amount to be:			refunded	\$		charged	\$	<p><b>CALCULATIONS PTO USE ONLY</b></p>	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE																																						
Total Claims	48 - 20 =	28	X \$18.00	\$	504.00																																				
Independent Claims	1 - 3 =	0	X \$84.00	\$	0																																				
MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes			+ \$280.00	\$	280.00																																				
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	1804.00																																				
	Amount to be:																																								
	refunded	\$																																							
	charged	\$																																							

a. ☒ A check in the amount of \$ **1804.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account. No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees.  
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
 overpayment to Deposit Account No. 02-2448.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

Send all correspondence to:  
**Birch, Stewart, Kolasch & Birch, LLP or Customer No. 2292**  
**P.O. Box 747**  
**Falls Church, VA 22040-0747**  
**(703) 205-8000**

**Date: April 3, 2002**

By Gerald M. Murphy, Jr.  
 Gerald M. Murphy, Jr., #28,977

/cqc

PATENT  
0760-0305P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: HIEI, Yukoh et al.  
Int'l. Appl. No.: PCT/JP00/05214  
Appl. No.: New Group:  
Filed: April 3, 2002 Examiner:  
For: METHOD FOR PROMOTING EFFICIENCY OF  
GENE TRANSFER INTO PLANT CELLS

PRELIMINARY AMENDMENT

**BOX PATENT APPLICATION**

Assistant Commissioner for Patents  
Washington, DC 20231

April 3, 2002

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

**AMENDMENTS**

**IN THE SPECIFICATION:**

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/JP00/05214 which has an International filing date of August 3, 2000, which designated the United States of America.--

Docket No. 0760-0305P

**IN THE CLAIMS:**

Please amend the claims as follows:

6. (Amended) The method according to claim 1 or 2, wherein the heat treatment is carried out for 5 seconds to 24 hours.

8. (Amended) The method according to claim 1 or 2, wherein the centrifugation is carried out under a centrifugal acceleration of 100G to 250,000G.

11. (Amended) The method according to claim 1 or 2, wherein said centrifugation acceleration of 1000G to 150,000G.

12. (Amended) A method for preparing a plant characterized by using the method according to claim 1 or 2.

13. (Amended) Plant cells, plant tissue or plant prepared by the method according to claim 1 or 2.

14. (Amended) The method according to claim 1 or 2, wherein said plant cells or plant tissue used are(is) originated from an angiosperm.

16. (Amended) Angiosperm cells, angiosperm tissue or angiosperm prepared by the method according to claim 14.

Docket No. 0760-0305P

19. (Amended) The monocotyledon cells, monocotyledon tissue or monocotyledon prepared by the method according to claim 17.

22. (Amended) The cells of the plant belonging to family Gramineae, the tissue of the plant belonging to family Gramineae, or the plant belonging to family Gramineae prepared by the method according to claim 20.

25. (Amended) Rice cells, rice tissue, rice, maize cells, maize tissue or maize prepared by the method according to claim 23.

Docket No. 0760-0305P

**REMARKS**

The specification has been amended to provide a cross-reference to the previously filed International Application.


The amendment to the claims is merely to correct improper multiple dependencies and to place the application into better form for examination. Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly solicited.

Attached hereto is a marked-up version of the changes made to the application by this Amendment.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By   
Gerald M. Murphy, Jr., #28,977

P.O. Box 747  
Falls Church, VA 22040-0747  
(703) 205-8000

GMM/cqc  
0760-0305P

Attachment: VERSION WITH MARKINGS TO SHOW CHANGES MADE

(Rev. 02/21/02)



Docket No. 0760-0305P

16. (Amended) Angiosperm cells, angiosperm tissue or angiosperm prepared by the method according to claim 14[ or 15].

19. (Amended) The monocotyledon cells, monocotyledon tissue or monocotyledon prepared by the method according to claim 17[ or 18].

22. (Amended) The cells of the plant belonging to family Gramineae, the tissue of the plant belonging to family Gramineae, or the plant belonging to family Gramineae prepared by the method according to claim 20[ or 21].

25. (Amended) Rice cells, rice tissue, rice, maize cells, maize tissue or maize prepared by the method according to claim 23[ or 24].

(Rev. 11/13/01)



4/pvt  
1

# SPECIFICATION

## Method for Promoting Efficiency of Gene Transfer into Plant Cells

### Technical Field

The present invention relates to a method for promoting efficiency of gene transfer into plant cells.

### Background Art

The method for transformation using *Agrobacterium* has a number of excellent features including, in general, the high efficiency, the small number of copies of the transferred gene, the feature that the gene may be transferred without fragmenting a specific region called T-DNA, and the feature that the frequency of mutation occurred during cultivation is low because transformants may be obtained by cultivation for a short period of time. Therefore, the method is widely used as the most useful method for transforming various plants.

Although the *Agrobacterium* method is an extremely excellent method for transforming plants, whether the transformation is successful or not and the transformation efficiency largely varies depending on the plant species, genotype and the plant tissue used (Potrykus et al. 1998 (Reference (36))). That is, there are species with which the transformation has not been successful, and species with which the transformation may be attained only with limited varieties. Further, there are species with which the tissue to be used is limited so that a large amount of materials cannot be treated. To prepare a practical variety by genetic recombination(genetic engineering), it is necessary to prepare a large number of transformed plants and to select the line having the desired character therefrom. However, at present, the type of plants with which a large number of transformed plants may be prepared for this purpose is limited. Thus, to develop an improved method by which this problem may be overcome is strongly demanded.

Although the method for transformation via *Agrobacterium* differs in the

On the other hand, studies for changing the plant tissue before infection of *Agrobacterium* to a physiological state in which the genes are likely to be transferred have been scarcely made. If the physiological state of the tissue can be changed to such a physiological state by a simple treatment, the method is very useful, and it is expected that, in addition to the promotion of the transformation efficiency, transformation may be attained for the species or genotypes with which transformation has been hitherto difficult, that is a prominent effect. Known studies about pretreatment of plant tissue include particle gun treatment (Bidney et al., 1992 (Reference (6))) and ultrasonication treatment (Trick H. N. et al., 1997 (Reference (40))). Both of these methods aim at promoting invasion of bacteria into the plant tissue by physically injuring the tissue, so as to increase the number of plant cells infected. However, these methods are nothing more than developments of the leaf disk method (Horsch et al., 1985 (Reference (19))) and not treatments based on novel concepts. The degree of effectiveness and universality of the methods have not been clarified, and they are not used as general methods.

### Disclosure of the Invention

The present inventors intensively studied to discover that in the gene transfer method using *Agrobacterium*, the gene transfer efficiency may be significantly promoted by heating and centrifuging the plant cells or plant tissue subjected to the gene transfer, thereby completing the present invention.

By the present invention, a method for promoting efficiency of gene transfer into plant cells, by which gene transfer can be attained simply with a higher efficiency than the conventional gene transfer by *Agrobacterium* method, without injuring the tissue, has been provided. The method of the present invention may be applied to both monocotyledons and dicotyledons. Further, the plants such as creeping bent grass which could not be hitherto transformed by *Agrobacterium* method can be transformed by the method of the present invention.

Fig. 1 is a drawing for showing a method for constructing pTOK233 which is an example of super-binary vectors, that may preferably be employed in the present invention.

Fig. 3 is a schematic view for showing the intermediate vector system and binary vector system which are major two vector systems of bacteria belonging to genus *Agrobacterium*.

Fig. 4 is a schematic view showing two binary vector systems derived from super virulent strain A281 of *Agrobacterium tumefaciens*.

In the above drawings, the following reference symbols denote the following meanings.

5 BL: left border sequence of T-DNA of bacteria belonging to genus

*Agrobacterium*

BR: right border sequence of T-DNA of bacteria belonging to genus  
*Agrobacterium*

TC: tetracycline resistance gene

10 SP: spectinomycin resistance gene

IG: intron GUS gene

HPT: hygromycin resistance gene

NPT: kanamycin-resistance gene

K: restriction enzyme *Kpn* I site

15 H: restriction enzyme *Hind* III site

Amp<sup>r</sup>: ampicillin resistance gene

BAR: bar gene

COS, cos: COS site of  $\lambda$  phage

ORI, ori: replication origin of ColE1

20 P35S: CaMV 35S promoter

Tnos: terminator of nopaline synthetase gene

virB: the *virB* gene in the virulence region of Ti plasmid pTiBo542  
contained in *Agrobacterium tumefaciens* A281

25 virC: the *virC* gene in the virulence region of Ti plasmid pTiBo542  
contained in *Agrobacterium tumefaciens* A281

virG: the *virG* gene in the virulence region of Ti plasmid pTiBo542  
contained in *Agrobacterium tumefaciens* A281

S Vir: entire *vir* region of Ti plasmid pTiBo542 of super virulent bacteria belonging to genus *Agrobacterium*

### Best Mode for Carrying out the Invention

The conditions of the heat treatment may appropriately be selected depending on the type of the plant used and the like, and may usually be carried out at a temperature of 30°C to 60°C, preferably 33°C to 55°C, more preferably 37°C to 52°C. The time of the heat treatment may appropriately be selected depending on the heating temperature, type of the plant used, the type of the cells or tissue to be heat-treated and so on, and is usually 5 seconds to 24 hours. When the heating temperature is high, the efficiency of transferring genes may be significantly promoted even if the time of the heat treatment is short. For example, when the heating temperature is 60°C, heat treatment for about 5 seconds may significantly promote the efficiency of gene transfer. On the other hand, when the heating

temperature of 30°C to 60°C, preferably 33°C to 55°C, more preferably 37°C to 52°C. The time of the heat treatment may appropriately be selected depending on the heating temperature, type of the plant used, the type of the cells or tissue to be heat-treated and so on, and is usually 5 seconds to 24 hours. When the heating temperature is high, the efficiency of transferring genes may be significantly promoted even if the time of the heat treatment is short. For example, when the heating temperature is 60°C, heat treatment for about 5 seconds may significantly promote the efficiency of gene transfer. On the other hand, when the heating

temperature is as low as about 34°C, the efficiency of gene transfer may be promoted by heat treatment for several tens of hours. In most cases, particularly preferred heating conditions are 37°C to 52°C for 1 minute to 24 hours, and the appropriate heating conditions for the particular plant cells or tissue may be easily selected by a routine experiment. By heating the plant cells or plant tissue at a temperature not lower than 55°C for a long time, the plant cells may be damaged and the efficiency of transformation may be decreased. Therefore, when the heating temperature is not lower than 55°C, the heating time is preferably short, for example, not longer than 3 minutes, preferably not longer than 1 minute so as to avoid damaging of the plant cells.

The conditions for centrifugation may appropriately be selected depending on the type of the plant used and the like, and may usually be carried out under a centrifugation acceleration of 100G to 250,000G, preferably 500G to 200,000G, more preferably 1000G to 150,000G. The time for centrifugation may appropriately be selected depending on the centrifugal acceleration, type of the plant used and so on, and is usually and preferably not less than one second. There is no upper limit of the centrifugation time, and about 10 minutes may usually be sufficient for attaining the object of the centrifugation. When the centrifugal acceleration is large, the efficiency of transferring genes may be significantly promoted even if the centrifugation time is very short, for example, 1 second or less. On the other hand, when the centrifugal acceleration is small, the efficiency of transferring genes may be significantly promoted by conducting the centrifugation for a long time. In most cases, especially preferred centrifugation conditions are about 500G to 200,000G, especially 1000G to 150,000G for about 1 second to 2 hours, and the appropriate centrifugation conditions for the particular plant cells or tissue may be easily selected by a routine experiment.

The method of the present invention is characterized by using the plant cells

or plant tissue which were(was) heated and centrifuged, or by contacting the plant cells or plant tissue with a bacterium belonging to the genus *Agrobacterium* while conducting the heat treatment and/or centrifugation, and as the method for gene transfer or transformation *per se* using the bacterium belonging to the genus *Agrobacterium*, a well-known method may be applied as it is.

The method for gene transfer or transformation *per se* into plants using a bacterium belonging to the genus *Agrobacterium* is well-known in the art and is widely used.

It is known for a long time that a soil bacterium *Agrobacterium* (*Agrobacterium tumefaciens*) causes crown gall disease in a number of dicotyledons. In 1970s, it was discovered that Ti plasmid concerns the virulence, and that the T-DNA which is a part of Ti plasmid is incorporated into the plant genome. Thereafter, it was proved that the T-DNA contains genes participating in synthesis of hormones (cytokinins and auxins) required for induction of tumor, and that the genes are expressed in plants in spite of the fact that the genes are bacterial genes. A group of genes existing in the virulence region (*vir* region) in the Ti plasmid is required for the excision of T-DNA and its transfer to plants, and the border sequences existing at the both ends of the T-DNA are necessary for the T-DNA to be excised. *Agrobacterium rhizogenes* which is another bacterium belonging to the genus *Agrobacterium* has a similar system on the Ri plasmid (Figs. 3 and 4).

Since T-DNA is incorporated into the plant genome by infection of *Agrobacterium*, it was expected that a desired gene may be incorporated into the plant genome by inserting the desired gene in the T-DNA. However, since Ti plasmid is as large as not less than 190 kb, it was difficult to insert a gene into the T-DNA by a standard technique of genetic engineering. Thus, a method for transferring a foreign gene into the T-DNA was developed.

First, disarmed strains such as LBA4404 (Hoekema et al., 1983 (Reference

(14))), C58C1(pGV3850) (Zambryski et al., 1983 (Reference (44))), and GV3Ti11SE (Fraley et al., 1985 (Reference (10))), that have tumorigenic Ti plasmids from which hormone synthetase genes were eliminated, were prepared (Fig. 3). Two methods employing such a strain, that is, a method by which a desired gene is transferred into the Ti plasmid of *Agrobacterium*, and a method by which a T-DNA having a desired gene is transferred into *Agrobacterium*, were developed. One of these methods is the so called intermediate vector method (Fraley et al., 1985 (Reference (10)); Fraley et al., 1983 (Reference (11)); Zambryski et al., 1983 (Reference (44)), Japanese Laid-open Patent Application (Kokai) No. 59-140885 (EP116718)). In this method, an intermediate vector which is easy to handle by genetic manipulation techniques, in which a desired gene may be inserted, and which can be replicated in *E. coli* is transferred into the T-DNA in the disarmed type Ti plasmid of *Agrobacterium* by triparental mating (Ditta et al., 1980 (Reference (9))). Another method is the so called binary vector method (Fig. 3), which is based on the fact that although the *vir* region is necessary for the T-DNA to be incorporated into plants, it is not necessary that the T-DNA and the *vir* region exist in the same plasmid ((Hoekema et al., 1983 . Reference (14)). The *vir* region contains *virA*, *virB*, *virC*, *virD*, *virE* and *virG* (Plant Biotechnology Encyclopedia (Enterprise Co., Ltd. (1989)), and the *vir* region is defined as those containing all of *virA*, *virB*, *virC*, *virD*, *virE* and *virG*. Thus, the binary vector is a small plasmid which is replicable in both *Agrobacterium* and *E. coli*, and this plasmid is transferred into *Agrobacterium* having a disarmed type Ti plasmid. The transferred of the binary vector into *Agrobacterium* may be carried out by electroporation method, triparental mating or the like). Binary vector includes pBIN19 (Bevan, 1984 (Reference (5))), pBI121 (Jefferson, 1987 (Reference (21))), pGA482 (An et al., 1988 (Reference (2))), Japanese Laid-open Patent Application (Kokai) No. 60-70080 (EP120516)), and a number of new binary vectors have been constructed based on these vectors. In the system of Ri



plasmid, similar vectors have been constructed and are used for transformation.

*Agrobacterium* A281 (Watson et al., 1975 (Reference (42))) is a super-virulent strain, whose host spectrum is wide and whose efficiency of transformation is higher than other strains (Hood et al., 1987(Reference (15)); Komari, 1989 (Reference (23))). This feature is brought about by a Ti plasmid pTiBo542 contained in A281 (Hood et al., 1984 (Reference (18)); Jin et al., 1987 (Reference (22)); Komari et al., 1986 (Reference (26))).

Two new systems using pTiBo542 has been developed. One system utilizes strains EHA101 (Hood et al., 1986, Reference (17)) and EHA105 (Hood et al., 1993, Reference (16)) containing a Ti plasmid which is a disarmed type of pTiBo542. By applying these strains to the above-mentioned binary vector system, a system having a high efficiency of transformation was achieved, which is widely used for transformation of various plants. Another system is "super-binary" vector system (Hiei et al., 1994 (Reference (13)); Ishida et al., 1996 (Reference (20)); Komari et al., 1999 (Reference (28)), WO94/00977, WO95/06722) (Fig. 4). Since this system comprises a disarmed type Ti plasmid having the *vir* region (*virA*, *virB*, *virC*, *virD*, *virE* and *virG*) (each of these may also be hereinafter referred to as "*vir* fragment region") and a plasmid having T-DNA, this is a kind of the binary vector system. However, it is different from the binary vector in that a super-binary vector (Komari, 1990a (Reference (24))) in which a *vir* region fragment (preferably a fragment containing at least *virB* or *virG*, more preferably a fragment at least containing *virB* and *virG*) substantially lacking at least one of the fragments of *vir* region is incorporated into the plasmid having the T-DNA, i.e., the binary vector. To transfer a T-DNA region into which a desired gene has been inserted into an *Agrobacterium* having the super-binary vector, homologous recombination via the triparental mating method may be employed as an easy method (Komari et al., 1996 (Reference (27))). It has been proved that the super-binary vector gives much higher transformation

In the method of the present invention, the host bacterium belonging to the genus *Agrobacterium* is not restricted, and *Agrobacterium tumefaciens* (e.g., the above-described *Agrobacterium tumefaciens* LBA4404 (Hoekema et al., 1983 (Reference (14))) and EHA101 (Hood et al., 1986 (Reference (17))) may preferably be employed.

The desired gene to be transferred into the plant may be inserted into a restriction site in the T-DNA region of the above-described plasmid by a conventional method, and the *Agrobacterium* into which the desired gene was

The desired gene to be transferred into the plant may be inserted into a restriction site in the T-DNA region of the above-described plasmid by a conventional method, and the *Agrobacterium* into which the desired gene was

incorporated may be selected based on an appropriate selection marker such as a drug resistant gene against a drug such as kanamycin or paromomycin. In cases where the plasmid is large and has a number of restriction sites, it is not always easy to insert the desired DNA into the T-DNA region by an ordinary subcloning method.

5 In such a case, the desired DNA may be inserted by the triparental mating method utilizing the homologous recombination in the cell of the bacterium belonging to the genus *Agrobacterium*.

Transfer of the plasmid into a bacterium belonging to the genus *Agrobacterium* such as *Agrobacterium tumefaciens* may be carried out by a known method including the above-mentioned triparental mating method, electroporation method, electroinjection method and chemical treatments with PEG or the like.

10 The gene which is to be transferred into the plant is, in principle, arranged between the left and right border sequences of the T-DNA as in the conventional method. However, since the plasmid is annular, the plasmid may contain only one border sequence. Alternatively, in cases where a plurality of genes are to be arranged at different sites, the plasmid may contain three or more border sequences. Alternatively, arrangement of the desired plasmid in the Ti or Ri plasmid may be performed in the cell of the bacterium belonging to the genus *Agrobacterium*, or the desired gene may be arranged in another plasmid. Further, the desired gene may be arranged in a plurality of types of plasmids.

20 Transfer of a gene into the plant cells via a bacterium belonging to the genus *Agrobacterium* may be attained by simply making the plant cells or plant tissue contact the bacterium belonging to the genus *Agrobacterium*. For example, a cell suspension of the bacterium belonging to the genus *Agrobacterium* having a population density of about  $10^6$  to  $10^{11}$  cells/ml is prepared, and the plant cells or the plant tissue are(is) immersed in the suspension for about 3 to 10 minutes, followed by co-culturing the resultant on a solid medium for several days, thereby attaining the

As the sample variety, Asanohikari which is a variety of Japonica rice was employed, and immature embryo was used as the material. Sample immature embryos were collected from immature seeds at 1 to 2 weeks after flowering and prepared by the method of Hiei, Y. et al (Reference (13)). That is, glumes of

immature seeds at 7 to 12 days after flowering were removed and the seeds were sterilized with 70% ethanol for 30 seconds and with 1% aqueous sodium hypochlorite solution for 10 minutes. Thereafter, immature embryos were excised and used as the samples. Calli derived from immature embryos were obtained by culturing the immature embryos on 2N6 medium (Hiei et al. 1994 (Reference (13)) (inorganic salts and vitamins of N6 (Chu C. C. 1978 (Reference (8), 1 g/l casamino acid, 2 mg/l 2,4-D) for two weeks.

As the *Agrobacterium* strains and plasmid vectors, LBA4404(pIG121Hm) (Hiei et al., 1994 (Reference (13))), LBA4404(pNB131) (see Fig. 2), and LBA4404(pTOK233) (Hiei et al., 1994 (Reference (13))) were used.

Construction of pNB131 was carried out as follows: After transferring pSB31 (Ishida Y, 1996 (Reference (20))) into *E. coli* LE392, pSB31 was transferred to *Agrobacterium* LBA4404 containing pNB1 (Komari T et al., 1996 (Reference (27))) by triparental mating method (Ditta G, 1980 (Reference (9))). By homologous recombination between pNB1 and pSB1 in the cell of *Agrobacterium*, pNB131 was obtained.

The T-DNA region of pIG121Hm contain a kanamycin resistant (nptII) gene driven by nos promoter, a hygromycin resistant (hpt) gene driven by 35S promoter of cauliflower mosaic virus (CaMV), and a GUS gene driven by the 35S promoter, which GUS gene contains introns of the catalase gene of castor bean (Ohta, S. et al., 1990 (Reference (33))).

The T-DNA region of pNB131 contains a bar gene driven by 35S promoter, and a GUS gene driven by the 35S promoter, which GUS gene contains introns (described above).

The T-DNA region of pTOK233 contains an nptII gene driven by nos promoter, an hpt gene driven by 35S promoter, and a GUS gene driven by the 35S promoter, which GUS gene contains introns (described above). The plasmid

## (2) Heat Treatment

### (3) Centrifugation Treatment

#### (4) Infection and Co-culturing

After the heat treatment or centrifugation treatment, or combination of these treatments, the sterilized water in each tube was removed and suspension of *Agrobacterium* was added, followed by stirring the mixture with a vortex mixer for 5 to 30 seconds. Preparation of the suspension of the bacterium was carried out by Hiei Y. et al. (Reference (13)). That is, colonies of *Agrobacterium* cultured on AB medium (Chilton, M-D et al., 1974 (Reference (7))) were collected with a platinum loop and the collected bacteria were suspended in modified AA medium (AA major inorganic salts, AA amino acids and AA vitamins (Toriyama K. et al., 1985 (Reference (39))), MS minor salts (Murashige, T et al., 1962 (Reference (32))), 1.0 g/l casamino acid, 100  $\mu$ M acetosyringone, 0.2 M sucrose, 0.2 M glucose). The population density of the bacterial cells in the suspension was adjusted to about 0.3 to  $1 \times 10^9$  cfu/ml. After leaving the mixture of immature embryos and the suspension of *Agrobacterium* to stand at room temperature for about 5 minutes, the immature embryos were plated on a medium for co-culturing. As the medium for co-culturing 2N6-AS medium (Hiei et al. 1994 (Reference (13)) containing 8 g/l agarose as a

#### (5) Selection of Transformed Cells (Japonica rice)

## (6) Results

### The results of the transient expression of the GUS gene after the heat

treatment and/or the centrifugation, and after the co-culturing with LBA4404(pIG121Hm) and LBA4404(pNB131) are shown in Tables 1 and 2. By carrying out the heat treatment or centrifugation treatment, the area in which GUS was expressed was clearly larger than the non-treated group, so that gene transfer occurred at a higher frequency. Further, by combining the heat treatment and the centrifugation treatment, the frequency was further increased.

The results of selection of the transformed calli obtained by culturing the rice immature embryos on the medium containing the selection drugs, after the co-culturing with *Agrobacterium* are shown in Tables 3, 4 and 5. The efficiency of obtaining transformed calli which were resistant to drug and which showed uniform expression of GUS gene was prominently increased by carrying out the heat treatment or the centrifugation treatment. Further, by combining the heat treatment and the centrifugation treatment, the efficiency of transformation was higher than in the cases where only one of these treatments was performed (Tables 3, 4 and 5). Thus, it was proved that by subjecting rice embryo to combination of the heat treatment and centrifugation treatment, efficiency of transformation was further promoted when compared with the cases where only one of these treatments was performed.

Further, it was confirmed that in cases where the efficiency of gene transfer is low by the centrifugation treatment alone due to the variety or the like, the efficiency of gene transfer was prominently increased by co-employing the heat treatment. Further, it was also confirmed that by setting the temperature of the centrifuge at about 40°C when the centrifugation is carried out, the centrifugation treatment and the heat treatment may be carried out simultaneously, and similar effect as the above-described combination of the treatments is obtained.

Hiei et al. (1994 (Reference (13))) reported that transformation may be attained with a relatively high efficiency using calli of rice. Aldemita RR et al.



1996 (Reference (1))) reported a case of transformation using rice immature embryo. To more effectively and more stably carry out these transformation methods, the above-described combined treatment method is very effective. Especially, although the quality of immature embryo is likely varied depending on the environment of culturing so that it is not easy to always obtain immature embryo suited for transformation, it may be possible to keep high efficiency of transformation by subjecting the immature embryo to the combined treatments. Hiei et al. (1994) (Reference (13)) showed that a super-binary vector having a high transformation ability promotes the efficiency of transformation of rice. According to Aldemita RR et al. 1996 (Reference (1))), transformants were obtained only in the test using LBA4404(pTOK233) containing a super-binary vector. By the combined treatment method according to the present invention, even when an ordinary binary vector is used, a high efficiency of transformation is attained, which is comparable to or even higher than that attained in the transformation using a super-binary vector. Further, by employing both the super-binary vector and the combined treatment method, the efficiency may be even more promoted. Still further, it is expected that transformants may be obtained by employing the combined treatment method for the varieties with which a transformant has not hitherto been obtained.

Table 1 Heat/Centrifugation Treatments and Transient Expression of GUS Gene in Scutella of Immature Embryos (Variety: Asanohikari)

Treatment Temperature (Treatment Time)	Centrifugal Acceleration (Treatment Time)	Number of Sample Immature Embryos	Number of Immature Embryos						
			Percentage of GUS-Expressed Area in Surface of Scutellum (%)						
			0	0-1	1-10	10-20	20-50	50-80	80-100
-	-	20	3	8	8	1	0	0	0
46°C (5 min)	-	20	1	6	7	4	2	0	0
-	20,000G (30 min)	20	0	1	4	7	7	1	0
46°C (5 min)	20,000G (30 min)	20	0	0	0	2	9	8	1

Sample Strain: LBA4404(pIG121Hm); Duration of Co-culturing: 5 days

Table 2 Heat/Centrifugation Treatments and Transient Expression of GUS

Gene in Scutella of Immature Embryos (Variety: Asanohikari)

Treatment Temperature (Treatment Time)	Centrifugal Acceleration (Treatment Time)	Number of Sample Immature Embryos	Number of Immature Embryos						
			Percentage of GUS-Expressed Area in Surface of Scutellum (%)						
			0	0-1	1-10	10-20	20-50	50-80	80-100
-	-	20	3	13	4	0	0	0	0
46°C (5 min)	-	20	0	0	10	7	3	0	0
-	20,000G (30 min)	20	0	0	3	9	8	0	0
46°C (5 min)	20,000G (30 min)	20	0	0	0	3	14	3	0

Sample Strain: LBA4404(pNB131); Duration of Co-culturing: 6 days

5

Table 3 Heat/Centrifugation Treatments and Efficiency of Selection of

Transformed Calli (Variety: Asanohikari)

Treatment Temperature (Treatment Time)	Centrifugal Acceleration (Treatment Time)	Number of Sample Sections of Immature Embryo (A)	Number of Hm-resistant GUS-Positive Calli (B)	B/A(%)
-	-	50	6	12.0
46°C (5 min)	-	51	15	29.4
-	20,000G (30 min)	51	29	56.9
46°C (5 min)	20,000G (30 min)	46	29	63.0

Sample Strain: LBA4404(pIG121Hm); Duration of Co-culturing: 5 days, Hm: 100

mg/l hygromycin

10

Table 4 Heat/Centrifugation Treatments and Efficiency of Selection of Transformed Calli (Variety: Asanohikari)

Treatment Temperature (Treatment Time)	Centrifugal Acceleration (Treatment Time)	Number of Sample Sections of Immature Embryo (A)	Number of Hm-resistant GUS-Positive Calli (B)	B/A(%)
-	-	60	7	11.7
46°C (5 min)	-	60	9	15.0
-	20,000G (1min)	60	48	80.0
-	20,000G (60 min)	60	48	80.0
46°C, 5 min	20,000G ( 1 min)	60	51	85.0
46°C, 5 min	20,000G (60 min)	60	51	85.0

Sample Strain: LBA4404(pIG121Hm); Duration of Co-culturing: 6 days, Hm: 100 mg/l hygromycin

5

Table 5 Heat/Centrifugation Treatments and Efficiency of Selection of Transformed Calli (Variety: Asanohikari)

Treatment Temperature (Treatment Time)	Centrifugal Acceleration (Treatment Time)	Number of Sample Sections of Immature Embryo (A)	Number of Hm-resistant GUS-Positive Calli (B)	B/A(%)
-	-	62	18	29.0
46°C (5 min)	-	64	32	52.5
-	20,000G (30 min)	60	39	65.0
46°C, 5 min	20,000G (30 min)	60	41	68.3

Sample Strain: LBA4404(pNB131); Duration of Co-culturing: 6 days, PPT: 10 mg/l phosphinothricin

## 10 Example 2

Immature embryos of maize with a size of about 1.2 mm (variety: A188, obtained from National Institute of Agrobiological Resources, The Ministry of Agriculture, Forestry and Fisheries) were aseptically collected and placed in a tube containing 2 ml of LS-inf liquid medium. After washing the embryos once with the same medium, 2.0 ml of fresh medium was added. Heat treatment was carried out by immersing the tube in a water bath at 46°C for 3 minutes. Centrifugation

15

treatment was performed by centrifuging the tube with a cooling centrifuge at 20 KG, at 4°C for 30 minutes. Combined heat/centrifugation treatments were carried out by conducting the above-described heat treatment and then conducting the above-described centrifugation treatment. The control was left to stand at room temperature for the same period of time. After the treatments, the medium was removed, and 1.0 ml of a suspension of *Agrobacterium tumefaciens* LBA4404(pSB131)(Ishida et al. 1986 (Reference (20))) with a population density of about  $1 \times 10^9$  cfu/ml in LS-inf liquid medium containing 100  $\mu$ M acetosyringone rubber was added, followed by stirring the mixture with a vortex mixer for 30 seconds. After leaving the mixture to stand at room temperature for 5 minutes, the embryos were plated on LS-AS medium containing 10  $\mu$ M AgNO<sub>3</sub> such that the surface of each hypocotyl contacts the medium. After co-culturing in the dark at 25°C for 3 days, an aliquot of the immature embryos was sampled and the transient expression of the GUS gene was checked by the treatment with X-gluc. The plasmid pSB131 is a super-binary vector.

The immature embryos after the co-culturing were cultured on a medium containing phsphinothricin (PPT) and 10  $\mu$ M AgNO<sub>3</sub>, thereby carrying out selection of the transformed cells. The calli grown on the selection medium were placed on a regeneration medium containing PPT, regeneration of transformed plants was carried out. A portion of a leaf of each regenerated plant was excised and the transient expression of the GUS gene was checked by the treatment with X-gluc as in Example 1. The above-described medium and method for culturing were in accordance with Ishida, Y. et al. 1996 (Reference (20)).

The results of transient expression of GUS gene when LBA4404(pSB131) was infected to the embryos after the treatments are shown in Table 6. All of the used immature embryos including the non-treated control, expression of GUS gene was observed. However, the area in which the GUS was expressed was larger in the

embryos subjected to heat treatment or combination of the heat treatment and centrifugation treatment than in the control. Especially, in the group subjected to the combination of the heat treatment and centrifugation treatment, the number of the embryos which showed expression of GUS gene in large area in the surface of the scutellum of each embryo was the largest.

The results of transformation of the immature embryos infected with LBA4404(pSB131) are shown in Table 7. From the control immature embryos which were not subjected to the heat treatment, transformed plants were obtained at a efficiency of 10.7%. On the other hand, the efficiency of transformation of the immature embryos subjected to the centrifugation treatment at 20KG at 4°C for 30 minutes was 13.3%, so that the efficiency was higher than that of the control group. The efficiency of the transformation of the embryos subjected to the heat treatment was 20%, which is about the twice that of the non-treated group. Further, the efficiency of transformation of the embryos subjected to the combined heat treatment and centrifugation treatment was 29.6% which was about three times that of the control group.

From the results described above, it was proved that the transformation efficiency is promoted by subjecting the immature embryos as the starting materials to centrifugation treatment or heat treatment before the infection, the transformation efficiency is even more promoted by combining these treatments. From these, the possibility that the maize varieties (Ishida et al. 1996 (Reference (20))) other than A188, which could not be hitherto transformed by the conventional *Agrobacterium* method, may be transformed by the centrifugation treatment, was suggested.

Table 6 Influence on Efficiency of Gene Transfer by Treatments (Infected with LBA4404(pSB131) )

Treatment	Sample Immature Embryo	G U S			
		+++	++	+	-
Not Treated	9	0	3	6	0
Heat	9	1	7	1	0
Centrifugation	12	0	3	9	0
Heat and Centrifugation	17	5	9	3	0

Transient expression of GUS gene in immature embryos after co-culturing

5 Table 7 Influence on Efficiency of Gene Transfer by Treatments (LBA4404(pSB131) was transferred)

Treatment	Number of Sample Immature Embryos	PPT-resistant callus (%)	PPT-resistant plant (%)	GUS+ plant (%)
Not Treated	28	9 (32.1)	9 (32.1)	3 (10.7)
Heat	30	18 (60.0)	15 (50.0)	6 (20.0)
Centrifugation	30	14 (46.6)	9 (30.0)	4 (13.3)
Heat and Centrifugation	27	23 (85.2)	20 (74.1)	8 (29.6)

Both the number of calli and number of plants do not include clones.

### Example 3

10 Mature seeds of creeping bent grass (*Agrostis palustris* cv. *Pencross*, Yukijirushi Shubyo Co., Ltd.) were sterilized and placed on a medium (TG2 medium) containing MS inorganic salts, MS vitamins, 4 mg/l dicamba, 0.5 mg/l 6BA, 0.7 g/l proline, 0.5 g/l MES, 20 g/l sucrose and 3 g/l gelrite (pH 5.8), followed by culturing the seeds at 25°C in the dark. The derived calli were subcultured on the medium having the same composition to grow embryogenic calli. The obtained  
 15 embryogenic calli were transferred to liquid medium (TG2L) which had the same composition as TG2 except that it did not contain gelrite, and cultured under shaking at 25°C in the dark to obtain cells of suspension culture. The cells of suspension

culture on 3 to 4 days after the subculture were placed in a tube containing 2 ml of TG2L medium. After once washing the calli with the same medium, 2 ml of fresh medium was added. The tube was immersed in a water bath at 46°C for 5 minutes. After removing the medium and adding the same fresh liquid medium, the resultant

5 was centrifuged at 5000 rpm at 4°C for 10 minutes. The control was left to stand at room temperature. The medium was removed and 1.0 ml of a suspension of *Agrobacterium tumefaciens* LBA4404(pTOK233) (described above) in TG2-inf medium (the same composition as TG2 medium except that proline, MES and gelrite are removed and 48.46 g/l sucrose, and 36.04 g/l glucose were added) at a population

10 density of about  $1 \times 10^9$  cfu/ml was added, followed by stirring the resulting mixture for 30 seconds by a vortex mixer. After leaving the calli to stand at room temperature for 5 minutes, the calli were placed on a medium (TG2-AS medium) which was the TG2L medium supplemented with 10 g/l glucose, 100 µM acetosyringone, 4 g/l type I agarose (pH5.8), and cultured at 25°C for 3 days i the

15 dark. The cells were then washed three times with TG2L medium containing 250 mg/l cefotaxime and carbenicillin. The cells were then suspended in the same buffer and cultured under shaking at 25°C in the dark at 70 rpm. One week later, the cells were subcultured on a medium having the same composition except that 50 mg/l hygromycin was added. After culturing the cells for another week, an aliquot

20 was sampled and treated with X-gluc to check the expression of the GUS gene.

Expression of the GUS gene in the suspended cultured cells of creeping bent grass infected with LBA4404(pTOK233) is shown in Table 8. In the control group, only one cell cluster showed expression of GUS. In contrast, in cases where the heat treatment and centrifugation treatment, about 30% of the cell clusters showed

25 expression of GUS gene. Further, the area in which the GUS gene was expressed was larger in the cell clusters subjected to heat and centrifugation treatments than that of the control group.

The transformation of creeping bent grass was hitherto only attained by the direct transfer method, i.e., by particle gun (Zhong et al. 1993 (Reference (45)), Hartman et al. 1994 (Reference (12)), Xiao, L. et al., 1997 (Reference (43))) or by electroporation (Asano Y., 1994 (Reference (3)), Asano Y. et al. 1998 (Reference (4))), and successful transformation by *Agrobacterium* method has not been reported. Assuming that the cause of the difficulty in transformation of creeping bent grass by *Agrobacterium* method is the low efficiency of the gene transfer in the known methods as can be seen from this Example, the possibility to obtain a transformed plant by the combined heat and centrifugation treatments according to the present invention was suggested.

Table 8      Effect of Heat and Centrifugation Treatments on Efficiency of Gene Transfer into Suspended Cultured Cells of Creeping Bent Grass

Treatment	Number of Cell Clusters		
	Total Number	GUS+	GUS+ (%)
Heat and Centrifugation Treatment	79	23	29.1
Control	101	1	1.0

Expression of GUS gene was checked two weeks after co-culturing

## References

- (1) Aldemita RR, Hodges TK (1996) *Agrobacterium tumefaciens*-mediated transformation of japonica and indica rice varieties. *Planta* 199: 612-617
- (2) An, G., Evert, P.R., Mitra, A. and Ha, S.B. (1988) Binary vectors. In Gelvin, S.B. and Schilperoort, R.A. (eds.), *Plant Molecular Biology Manual A3*. Kluwer Academic Press, Dordrecht, pp. 1-19.
- (3) Asano, Y., Ugaki, M. (1994) Transgenic plants of *Agrostis alba* obtained by electroporation-mediated direct gene transfer into protoplasts. *Plant Cell Reports* 13:243-246.
- (4) Asano, Y., Ito, Y., Fukami, M., Sugiura, K., Fujiie, A. (1998) Herbicide-resistant



transgenic creeping bentgrass plants obtained by electroporation using an altered buffer. Plant Cell Reports 17:963-967.

(5) Bevan, M. (1984) Binary Agrobacterium vectors for plant transformation. Nucleic Acids Res., 12, 8711-8721.

5 (6) Bidney, D., Scelongo, C., Martich, J., Burrus, M., Sims, L., and Huffman G. (1992) Microprojectile bombardment of plant tissues increases transformation frequency by Agrobacterium tumefaciens. Plant Mol. Biol., 18, 301-313.

(7) Chilton, M-D., Currier, TC. Farrand, SK. Bendich, AJ. Gordon, MP. & Nester EW. (1974) Agrobacterium tumefaciens DNA and PS8 bacteriophage DNA not  
10 detected in crown gall tumors. Proc. Natl. Acad. Sci. USA, 71:3672-3676

(8) Chu, C. C., (1978) Proc. Symp. Plant Tissue Culture, Science Press Peking, pp.43-50

(9) Ditta, G., Stanfield, S., Corbin, D. and Helinski, D.R. (1980) Broad host range DNA cloning system for Gram-negative bacteria: Construction of gene bank of  
15 Rhizobium meliloti. Proc. Natl. Acad. Sci. USA, 77, 7347-7351.

(10) Fraley, R.T., Rogers, S.G., Horsch, R.B., Eicholtz, D.A. and Flick, J.S. (1985) The SEV system: a new disarmed Ti plasmid vector for plant transformation. Bio/technology, 3, 629-635.

(11) Fraley, R.T., Rogers, S.G., Horsch, R.B., Sanders, P.R., Flick, J.S., Adams, S.P.,  
20 Bittner, M.L., Brand, L.A., Fink, C.L., Fry, J.S., Galluppi, G.R., Goldberg, S.B., Hoffmann, N.L. and Woo, S.C. (1983) Expression of bacterial genes in plant cells. Proc Natl Acad Sci USA, 80, 4803-4807.

(12) Hartman, C.L., Lee, L., Day, P.R., Turner, N.E. (1994) Herbicide resistant turfgrass (*Agrostis palustris* Huds.) by biolistic transformation. Biotechnology  
25 12:919-923.

(13) Hiei, Y., Ohta, S., Komari, T. and Kumashiro, T. (1994) Efficient transformation of rice (*Oryza sativa* L.) mediated by Agrobacterium and sequence

analysis of the boundaries of the T-DNA. The Plant Journal, 6, 271-282.

(14) Hoekema, A., Hirsch, P.R., Hooykaas, P.J.J. and Schilperoort, R.A. (1983) A binary plant vector strategy based on separation of vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. Nature, 303, 179-180.

5 (15) Hood, E.E., Fraley, R.T. and Chilton, M.-D. (1987) Virulence of *Agrobacterium tumefaciens* strain A281 on legumes. Plant Physiol, 83, 529-534.

(16) Hood, E.E., Gelvin, S.B., Melchers, L.S. and Hoekema, A. (1993) New *Agrobacterium* helper plasmids for gene transfer to plants. Transgenic Res., 2, 208-218.

10 (17) Hood, E.E., Helmer, G.L., Fraley, R.T. and Chilton, M.-D. (1986) The hypervirulence of *Agrobacterium tumefaciens* A281 is encoded in a region of pTiBo542 outside of T-DNA. J. Bacteriol., 168, 1291-1301.

(18) Hood, E.E., Jen, G., Kayes, L., Kramer, J., Fraley, R.T. and Chilton, M.-D. (1984) Restriction endonuclease map of pTiBo542, a potential Ti-plasmid vector for genetic engineering of plants. Bio/technology, 2, 702-709.

15 (19) Horsch, R. B., Fry, J. E., Hoffmann, N. L., Eichholtz, D., Rpgers, S. G. and Fraley, R. T. (1985) A simple and general method for transferring genes into plants. Science 227, 1229-1231.

(20) Ishida, Y., Saito, H., Ohta, S., Hiei, Y., Komari, T. and Kumashiro, T. (1996) High efficiency transformation of maize (*Zea mays* L.) mediated by *Agrobacterium tumefaciens*. Nature Biotechnol, 14, 745-750.

(21) Jefferson, R.A. (1987) Assaying chimeric genes in plants: the GUS gene fusion system. Plant Mol. Biol. Rep., 5, 387-405.

20 (22) Jin, S., Komari, T., Gordon, M.P. and Nester, E.W. (1987) Genes responsible for the supervirulence phenotype of *Agrobacterium tumefaciens* A281. J. Bacteriol., 169, 4417-4425.

(23) Komari, T. (1989) Transformation of callus cultures of nine plant species

mediated by *Agrobacterium*. *Plant Sci.*, 60, 223-229.

(24) Komari, T. (1990a) Genetic characterization of double-flowered tobacco plant obtained in a transformation experiment. *Theor. Appl. Genet.*, 80, 167-171.

(25) Komari, T. (1990b) Transformation of cultured cells of *Chenopodium quinoa* by binary vectors that carry a fragment of DNA from the virulence region of pTiBo542. *Plant Cell Reports*, 9, 303-306.

(26) Komari, T., Halperin, W. and Nester, E.W. (1986) Physical and functional map of supervirulent *Agrobacterium tumefaciens* tumor-inducing plasmid pTiBo542. *J. Bacteriol.*, 166, 88-94.

(27) Komari, T., Hiei, Y., Saito, Y., Murai, N. and Kumashiro, T. (1996) Vectors carrying two separate T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. *Plant J.*, 10, 165-174.

(28) Komari, T. and Kubo, T. (1999) *Methods of Genetic Transformation: Agrobacterium tumefaciens*. In Vasil, I.K. (ed.) *Molecular improvement of cereal crops*. Kluwer Academic Publishers, Dordrecht, pp. 43-82.

(29) Li, H.-Q., Sautter, C., Potrykus, I. and Puonti-Kaerlas, J. (1996) Genetic transformation of cassava (*Manihot esculenta* Crantz). *Nature Biotechnol.*, 14, 736-740.

(30) Lindsey, K., Gallois, P. and Eady, C. (1991) Regeneration and transformation of sugarbeet by *Agrobacterium tumefaciens*. *Plant Tissue Culture Manual* B7:1-13. Kluwer Academic Publishers.

(31) McCormick, S. (1991) Transformation of tomato with *Agrobacterium tumefaciens*. *Plant Tissue Culture Manual* B6:1-9. Kluwer Academic Publishers.

(32) Murashige, T. and Skoog, F. (1962) *Physiol. Plant* 15:473-497.

(33) Ohta, S., Mita, S., Hattori, T., Namamura, K. (1990) Construction and expression in tobacco of a  $\beta$ -glucuronidase (GUS) reporter gene containing an intron

within the coding sequence. *Plant Cell Physiol.* 31: 805-813.

(34) Potrykus I., Harms, C. T. and Lorz, H. (1979) Callus formation from cell culture protoplasts of corn (*Zea mays* L.). *Theor. Appl. Genet.* 54:209-214.

(36) Potrykus, I., Bilang, R., Futterer, J., Sautter, C. and Schrott, M. (1998)  
5 *Agricultural Biotechnology*, NY:Mercel Dekker Inc. pp. 119-159.

(37) Rogers, S.G., Horsch, R.B. and Fraley, R. T. (1988) Gene transfer in plants: Production of transformed plants using Ti plasmid vectors. *Method for Plant Molecular Biology*, CA: Academic Press Inc. pp.423-436.

(38) Saito, Y., Komari, T., Masuta, C., Hayashi, Y., Kumashiro, T. and Takanami, Y.

10 (1992) Cucumber mosaic virus-tolerant transgenic tomato plants expressing a satellite RNA. *Theor. Appl. Genet.*, 83, 679-683.

(39) Toriyama, K. and Hinata, K. (1985) *Plant Sci.* 41:179-183

(40) Trick, H.N. and Finer, J.J. (1997) SAAT: sonication-assisted *Agrobacterium*-mediated transformation. *Transgenic Research* 6:329-336.

15 (41) Visser, R.G.F. (1991) Regeneration and transformation of potato by *Agrobacterium tumefaciens*. *Plant Tissue Culture Manual* B5:1-9. Kluwer Academic Publishers.

(42) Watson, B., Currier, T.C., Gordon, M.P., Chilton, M.-D. and Nester, E.W.  
(1975) Plasmid required for virulence of *Agrobacterium tumefaciens*. *J Bacteriol.*  
20 123, 255-264.

(43) Xiao, L., Ha, S.-B. (1997) Efficient selection and regeneration of creeping bentgrass transformants following particle bombardment. *Plant Cell reports* 16:874-878.

(44) Zambryski, P., Joos, H., Genetello, C., Leemans, J., Van Montagu, M. and  
25 Schell, J. (1983) Ti plasmid vector for the introduction of DNA into plant cells without alteration of their normal regeneration capacity. *EMBO J*, 2, 2143-2150.

(45) Zhong, H., Bolyard, M.G., Srinivasan, C., Sticklen, M.B. (1993) *Transgenic*

plants of turfgrass (*Agrostis palustris* Huds.) from microprojectile bombardment of embryogenic callus. Plant Cell Reports 13:1-6.

---

## CLAIMS

1. A method for promoting efficiency of gene transfer into plant cells by a bacterium belonging to genus *Agrobacterium*, comprising heating and centrifuging said plant cells or plant tissue.
2. The method according to claim 1, wherein said gene transfer is carried out after heating and centrifuging said plant cells or plant tissue.
3. The method according to claim 1 or 2, wherein the heat treatment is carried out at a temperature of 33°C to 60°C.
4. The method according to claim 3, wherein the heat treatment is carried out at a temperature of 35°C to 55°C.
5. The method according to claim 4, wherein the heat treatment is carried out at a temperature of 37°C to 52°C.
6. The method according to any one of claims 1 to 5, wherein the heat treatment is carried out for 5 seconds to 24 hours.
7. The method according to claim 1 or 2, wherein the heat treatment is carried out at a temperature of 37°C to 52°C for 1 minute to 24 hours.
8. The method according to any one of claims 1 to 7, wherein the centrifugation is carried out under a centrifugal acceleration of 100G to 250,000G.
9. The method according to claim 8, wherein said centrifugation is carried out under a centrifugal acceleration of 500G to 200,000G.
10. The method according to claim 9, wherein said centrifugation is carried out under a centrifugal acceleration of 1000G to 150,000G.
11. The method according to any one of claims 1 to 10, wherein said centrifugation is carried out for 1 second to 4 hours.
12. A method for preparing a plant characterized by using the method according to claim 1 to 11.
13. Plant cells, plant tissue or plant prepared by the method according to claims 1

to 12.

14. The method according to any one of claims 1 to 11, wherein said plant cells or plant tissue used are(is) originated from an angiosperm.

15. A method for preparing an angiosperm characterized by using the method according to claim 14.

16. Angiosperm cells, angiosperm tissue or angiosperm prepared by the method according to claim 14 or 15.

17. The method according to claim 14, wherein said plant cells or plant tissue used are(is) originated from a monocotyledon.

18. A method for preparing a monocotyledon characterized by using the method according to claim 17.

19. The monocotyledon cells, monocotyledon tissue or monocotyledon prepared by the method according to claim 17 or 18.

20. The method according to claim 17, wherein said plant cells or plant tissue are(is) originated from a plant belonging to family Gramineae.

21. A method for preparing a plant belonging to family Gramineae characterized by using the method according to claim 20.

22. The cells of the plant belonging to family Gramineae, the tissue of the plant belonging to family Gramineae, or the plant belonging to family Gramineae prepared by the method according to claim 20 or 21.

23. The method according to claim 20, wherein said plant cells or plant tissue are(is) of rice or maize.

24. A method for preparing rice or maize characterized by using the method according to claim 23.

25. Rice cells, rice tissue, rice, maize cells, maize tissue or maize prepared by the method according to claim 23 or 24.

A method for gene transfer by which higher efficiency for gene transfer than that by the conventional *Agrobacterium* method may be attained simply and without injuring the tissue is disclosed. According to the method of the present invention, the efficiency of gene transfer into plant cells by a bacterium belonging to genus *Agrobacterium* is promoted by accompanying heat treatment and centrifugation treatment of the plant cells or plant tissue.



1/4




Fig. 1

2/4




Fig. 2

3/4




Fig. 3

4/4




Fig. 4

BIRCH, STEWART, KOLASCH &amp; BIRCH, LLP

0760-0305P

P.O. Box 747 · Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 · Facsimile: (703) 205-8050COMBINED DECLARATION AND POWER OF ATTORNEY  
FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed, and for which a patent is sought on the invention entitled: "Method for Promoting Efficiency of Gene Transfer into Plant Cells"

the specification of which is attached hereto. If not attached hereto,  
the specification was filed on \_\_\_\_\_ as  
United States Application Number \_\_\_\_\_;  
and amended on \_\_\_\_\_ (if applicable) and/or  
the specification was filed on 08/03/00 as PCT  
International Application Number PCT/JP00/05214; and was  
amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representative or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)		Priority Claimed
<u>158026/99</u>	<u>Japan</u>	<u>06/04/99</u>
(Number)	(Country)	(Month/Day/Year Filed)
_____	_____	_____
(Number)	(Country)	(Month/Day/Year Filed)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional applications(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Country	Application Number	Date of Filing (Month/Day/Year)
---------	--------------------	---------------------------------

_____	_____	_____
-------	-------	-------

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

_____	_____	_____
(Application Number)	(Filing Date)	(Status - patented, pending, abandoned)
_____	_____	_____
(Application Number)	(Filing Date)	(Status - patented, pending, abandoned)

Attorney Docket No.

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart  
Joseph A. Kolasch  
Bernard L. Sweeney  
Charles Gorenstein  
Leonard R. Svensson  
Andrew D. Meikle  
Joe McKinney Muncy  
John W. Bailey  
Gary D. Yacura

(Reg. No. 21,066)  
(Reg. No. 22,463)  
(Reg. No. 24,448)  
(Reg. No. 29,271)  
(Reg. No. 30,330)  
(Reg. No. 32,868)  
(Reg. No. 32,334)  
(Reg. No. 32,881)  
(Reg. No. 35,416)

Terrell C. Birch  
James M. Slattery  
Michael K. Mutter  
Gerald M. Murphy, Jr.  
Terry L. Clark  
Marc S. Weiner  
Donald J. Daley  
John A. Castellano

(Reg. No. 19,382)  
(Reg. No. 28,380)  
(Reg. No. 29,680)  
(Reg. No. 28,977)  
(Reg. No. 32,644)  
(Reg. No. 32,181)  
(Reg. No. 34,313)  
(Reg. No. 35,094)

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP** or  
P.O. Box 747 · Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 · Facsimile: (703) 205-8050

**Customer No. 2292**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00

GIVEN NAME/FAMILY NAME <b>Yukoh HIEI</b>	INVENTOR'S SIGNATURE <i>Yukoh Hiei</i>	DATE* Apr. 9, 2002
Residence (City, State & Country) <b>Iwata-gun, Shizuoka, JAPAN JPX</b>		CITIZENSHIP <b>Japanese -</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <b>c/o Orynova K.K., 700, Higashibara, Toyoda-cho, Iwata-gun, Shizuoka, 438-080 2 JAPAN</b>		

2-00

GIVEN NAME/FAMILY NAME <b>Keisuke KASAOKA</b>	INVENTOR'S SIGNATURE <i>Keisuke Kasaka</i>	DATE* Apr. 9, 2002
Residence (City, State & Country) <b>Oyama-shi, Tochigi, JAPAN JPX</b>		CITIZENSHIP <b>Japanese -</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <b>c/o Japan Tobacco Inc., Leaf Tobacco Research Center, 1900, Idei, Ooaza, Oyama-shi, Tochigi, 323-0808 JAPAN</b>		

3-00

GIVEN NAME/FAMILY NAME <b>Yuji ISHIDA</b>	INVENTOR'S SIGNATURE <i>Yuji Ishida</i>	DATE* Apr. 9, 2002
Residence (City, State & Country) <b>Iwata-gun, Shizuoka, JAPAN JPX</b>		CITIZENSHIP <b>Japanese -</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <b>c/o Orynova K.K., 700, Higashibara, Toyoda-cho, Iwata-gun, Shizuoka, 438-0802 JAPAN</b>		

GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		

\*DATE OF SIGNATURE